

Intervention for Social Anxiety among University Students with a Solution-Focused Group Counseling Program

Mehmet Enes Sağarⁱ
Afyon Kocatepe University

Abstract

The aim of the research is to investigate the effect of a solution-focused group counseling program on reducing social anxiety of university students. The study group consists of 30 (16 female, 14 male) students who volunteered among 224 students who continue their education at Afyon Kocatepe University in the 2019-2020 academic year. The research had a 2x3 mixed pattern with experiment and control groups and pretest, posttest and follow-up measures. The Liebowitz Social Anxiety Scale (LSAS) was used as data collection tool. University students in the experimental group attended a 6-session solution-focused group counseling program about reducing social anxiety, while the control group did not participate in any study. After completing experimental processes, both groups underwent posttest and then follow-up measures at the end of 3 months. Data were analyzed with the two-way analysis of variance technique for mixed patterns. Results revealed that the solution-focused group counseling program was effective in reducing the social anxiety of university students and the efficacy continued during follow-up at the end of three months.

Keywords: Solution-Focused Group Counseling, Social Anxiety, Program, University, Student

DOI: 10.29329/ijpe.2021.346.20

ⁱ **Mehmet Enes Sağar**, Dr., Department of Child Care and Youth Services, Afyon Kocatepe University, Çay Vocational School, ORCID: 0000-0003-0941-5301

Email: mehmetenes15@gmail.com

INTRODUCTION

In order for the social beings of humans to maintain an adjusted, balanced and regular life during their whole lives, it is undoubtedly very important not to break social bonds with other people and to maintain relationships. In the youth period, abandoning their original social groups and beginning university life is a habitual life cycle. However, during this period, young people may encounter significant difficulties in terms of adjustment (Deci & Ryan, 2004). Facing many personal and academic difficulties during this process, young people may experience emotional distress during their experiences and this may make the adjustment process even more difficult (Bolsoni-Silva & Loureiro, 2014; Mustafa, Hamdan-Mansour, Hijazeen, Abed, Abdallah, El-Haija & Omari, 2014; Wintre & Yaffe, 2000). In addition to these difficulties encountered during university life, young people with less interaction may feel inadequate and remain distant from their social environment (Ateş & Gençdoğan, 2017; Elemo, 2019). This worrying situation for young people is qualified as social anxiety and is defined as “individuals avoiding one or more social situations due to experiencing pronounced forms of fear or anxiety in relation to exposure to possible investigation or negative assessment by others” (American Psychiatric Association, 2013). In other words, it is defined as the tendency of individuals to fear or avoid situations that they think are examined or evaluated by others (Rapee, 1995).

There is a need for increased studies investigating treatment and therapy approaches related to social anxiety problems among young people. In the relevant literature, there are studies investigating efficacy based on psychological counselling approaches to overcome this problem and meta-analyses reviewing these studies. Studies aimed at reducing social anxiety in the literature are generally in the form of group counseling. It has been observed that these studies are based on cognitive-behavioral approach (Aydn, Sütçü Tekinsav & Sorias, 2010; Abeditehrani, Dijk, Toghchi & Arntz, 2020; Fogarty, Hevey & McCarthy, 2019; Mercan, 2007; Tillfors, Andersson, Ekselius, Furmark, Lewenhaupt, Karlsson & Carlbring, 2011), behavioral counseling approach (Beidel, Turner & Morris, 2000) and reality theory (Palancı, 2014). A meta-analysis completed in 2013 revealed that cognitive behavioral psychological counseling, exposure and social skills, supported self-help, unsupported self-help and psychodynamic psychological counseling were effective to cope with social anxiety; however, individual cognitive behavioral psychological counseling was more effective (Mayo-Wilson, Dias, Mavranzouli, Kew, Clark, Ades & Pilling, 2014). Another meta-analysis study stated that exposure, cognitive restructuring techniques and social skills education and a combination of these techniques were effective; however, studies based on cognitive behavioral approaches had higher effect magnitude. This result indicates that the cognitive behavioral approach was effective on social anxiety (Gil, Carrillo & Meca, 2001). In line with this knowledge, it appears that studies about coping with social anxiety problems have focused on cognitive approaches. In terms of coping with social anxiety problems, it appears there are studies based on solution-focused approaches, especially in the international literature (Bajjesh, 2015; Elemo, 2019; Esmail, Alireza, Khadije, Esmail & Shima, 2019; George, 2008; Mahdiyar, Dadfarnia, Hadianfard & Rahimi, 2019). However, in the Turkish literature, it can be said that research based on a solution-focused psychological counseling approach to reducing social anxiety is quite inadequate. It is notable as an approach with efficacy in a short duration for this common problem among young people. In this context, the effectiveness of the solution-focused counseling approach as an alternative to other counseling approaches in dealing with social anxiety can also be tested.

Solution-focused counseling approach developed by Steve de Shazer and his team is accepted among counseling approaches in the whole world developed in the last fifty years in America. It is based on rules like “if it isn’t broken, don’t fix it”, “apply functional solution paths” and “don’t insist on trying dysfunctional solution paths, try different solutions”. People focus on solution paths instead of problems and assist in producing targets and solutions as experts in their own lives. Solution-focused counseling is a collaborative process that takes place between 4 and 6 sessions and puts the client at the center. The techniques and basic components of this approach are miracle question, exceptions, scaling questions, praise, homework, emphasizing strong aspects, searching for solutions

and determining targets (De Shazer, 1985; De Shazer & Berg, 1997; De Jong & Berg, 1998; Doğan, 1999; Kim & Franklin, 2009; Simon & Berg, 1997; Sklare, 1997). In the context of these techniques based on the solution-focused approach, it is significant how the solution-focused approach provides a new dimension as an alternative to other problem-focused approaches for university students coping with social anxiety. It is thought that there is a need for solution-oriented studies and programs that can be used functionally in the field of psychological counseling to intervene in the social anxiety of university students. Additionally, the need for university students to integrate their daily life activities with society in more regular, adjusted and balanced fashion makes this topic important. For these reasons, in this study, it is tried to find answers to the questions about to what extent the program based on the solution-focused group counseling approach will contribute to reducing the social anxiety of university students. Within this target framework, the answer to the following hypothesis was researched:

Hypothesis: When university students in the solution-focused group counseling group and control group are compared, there will be a significant reduction in posttest social anxiety scale points compared to pretest points and this reduction will continue during follow-up measures performed 3 months after the sessions end.

MATERIALS AND METHODS

Research Pattern

This research is a quasi-experimental study using 2 x 3 split-plot factorial (mixed) pattern. The first factor in this pattern is the independent process groups of the experiment and control groups, while the second factor represents the dependent variable of repeated measures at different times of the pretest, posttest and follow-up measurements (Büyüköztürk, 2011). Due to field studies, it is difficult to provide real experimental design conditions, especially to assign randomly to groups. Therefore, quasi-experimental design offers greater flexibility for many applied researches. Quasi-experimental designs involve the manipulation of one or more independent variables as in real experimental designs. Random assignment of study participants is not included. It is as useful as possible in applied research due to both creativity in the use of patterns and appropriate use of statistical processes and controls (Heppener, Wampold, & Kivlighan, 2008). Therefore, quasi-experimental design was preferred in the framework of this research.

Study Group

The study group was selected from volunteer university students who received low scores by completing the "Liebowitz Social Anxiety Scale-Turkish Form (LSAS)" scale tool at Afyon Kocatepe University. The research was completed with a total of 30 university students with 15 in the experiment group and 15 in the control group. The study groups consist of 8 female, 7 male and 15 students.

Data Collection Tools

With the aim of gathering data in the research, "Liebowitz Sosyal Anxiety Scale" was used.

Liebowitz Social Anxiety Scale-Turkish Form (LSAS): Liebowitz Social Anxiety Scale (LSAS) was developed by Heimberg, Horner, Juster, Safren, Brown, Schneier and Liebowitz (1999). This scale, adapted to Turkish by Soykan, Devrimci, Özgüven and Gençöz (2003), was prepared to determine the levels of fear and/or avoidance of individuals in performance situations with social interaction. The scale comprises 24 items with Likert-type responses from 1-4. Inter-assessor reliability of the scale was 0.96, test-repeat test reliability was 0.97 and internal consistency (Cronbach alpha coefficient) was 0.98 (Soykan, Devrimci, Özgüven & Gençöz, 2003).

Procedure

Ethics committee permission was obtained from Afyon Kocatepe University Social and Humanities Scientific Research and Publication Ethics Committee (Date of Decision: 06.02.2020; Meeting: 01; Number of Documents: E.6467). Then the LSAS was applied to 224 volunteer university students attending University with the aim of determining participants. Then, the 224 university students were ranked from lowest to highest for LSAS points. Thirty university students with high social anxiety points who agreed to participate in the study voluntarily as a result of preliminary interviews provided written consent and were randomly placed in experiment and control groups with the lottery method.

Before beginning sessions, the experiment and control groups had LSAS pretest applied. The independent group t test was performed to determine whether the experiment and control groups were equivalent and there was no significance for the difference between the pretest mean points for social anxiety ($t = 1.05, p > .05$) in the groups. In line with this, it may be stated that the experiment and control groups had equivalent social anxiety levels before the implementation. The independent group t test results for pretest social anxiety points of university students in the experiment and control groups are given in Table 1.

Table 1. Independent Group T test Results related to LSAS pretest points

Groups	N	Mean	Ss	t	P
Experimental Group	15	109.33	15.74	1.05	.301
Control Group	15	104.20	10.39		

Ss: Standard deviation, t: t test

Later the experimental group underwent 6 sessions of a solution-focused group counseling program lasting 120 minutes each week. There was no implementation for the control group. After completing sessions, participants in both groups were presented with the LSAS posttest. Three months after the posttest administration, later the same experiment and control groups had the follow-up test applied.

Solution-Focused Group Counseling Program

The general aim of this program was to allow university students to reduce social anxiety with a solution-focused perspective and live in a more-adjusted way. During development of the program, the domestic and international literature was investigated (Ateş & Gençdoğan, 2017; Bajesh, 2015; De Shazer, 1985; Doğan, 1999; Elemo, 2019; Mahdiyar, Dadfarnia, Hadianfard & Rahimi, 2019; Baygül, 2015; Sarı & Günaydın, 2015). This program focuses on small changes and exceptions and includes solution-focused techniques for clients to discover their strong aspects, and assist in searching for solutions and determining targets; to use scaling questions and coping techniques; to ask the miracle question and ensure a positive view of the future; to do homework; to compliment clients; and to encourage clients; in addition to activities like directed dream studies, finding positive stories, discussion and role-play. After the preparation and development stage of the solution-focused group counseling program, pilot applications were completed with a 10-person group to create the program framework. During the pilot study, inadequacies related to the program were identified and revisions were made and the program was made appropriate for the study group. During implementation, care was taken to complete group sessions on the determined date and durations.

The content summary of the solution-focused group counseling program developed in the context of this research is given in Table 2.

Solution-Focused Group Counseling Program

Table 2. Content Of Solution-Focused Counseling Program

Solution-Focused Group Counseling Program				
Session	Number of participants	Duration	Aim	Solution-focused techniques
1st session	15	120 minutes	Opening, group formation, ensuring solution-focused approach and awareness of coping sources	Variation Technique Before Session, Power Sources, Encouragement, Exceptions
2nd session	15	120 minutes	Ensuring clear determination of achievable aims	Miracle Question Technique
3rd session	15	120 minutes	Ensuring awareness of power sources and times without problems	Exceptions/Rare Situations Technique, Coping Questions Technique, Imagination Questions Technique
4th session	15	120 minutes	Ensuring awareness of methods used when coping with problems and times when successfully used	Exceptions/Rare Situations Technique, Coping Questions Technique
5th session	15	120 minutes	Ensuring an idea about what life will be like when problems are solved in the future	Scaling Questions Technique, Future-Focused Questions Technique, Crystal Ball Technique, Exceptions/Rare Situations Technique, Coping Questions Technique
6th session	15	120 minutes	Ensuring an assessment of the group process and ending this process	Exception, encouragement

Analysis of Data

Firstly, the distribution graph of data and the Shapiro-Wilks normality test were used to determine whether there were parametric values in the experiment and control groups in the research. Both groups were identified to show normal distribution. Analysis of data in line with this used the two-factor variance analysis technique for mixed patterns. Analysis of data was completed in a computer environment with the SPSS program and analyses assessed significance level at .05 (Büyüköztürk, 2011).

FINDINGS

Before testing the trial in the research, firstly the mean and standard deviations were calculated for points obtained on the LSAS by the experiment and control groups of university students before the implementation, after the implementation and 3 months after the implementation ended. The values are presented in Table 3.

Table 3. Mean and Standard Deviation of LSAS Pretest, Posttest and Follow-up Test Points for Experiment and Control Groups

Groups	n	Pretest		Posttest		Follow-up test	
		Mean	Standard deviation	Mean	Standard deviation	Mean	Standard deviation
Experimental Group	15	109.33	15.74	62.53	6.09	61.80	5.64
Control Group	15	104.20	10.39	104.93	10.91	101.66	12.95
Total	30	106.76	13.36	83.73	23.24	81.73	22.52

As observed in Table 3, participants in the experimental group were observed to have lower mean points on the LSAS posttest ($\bar{X}=62.53$) and 3-month follow-up test ($\bar{X}=61.80$) compared to the pretest ($\bar{X}=109.33$).

In terms of testing the trial in the research, the two-factor variance analysis technique for split-plot patterns was used to analyze whether the difference between mean points on the pretest, posttest and follow-up tests were significant or not. The obtained findings are presented in Table 4. The analysis results on Table 4 did not abide by the Mauchly's sphericity assumption ($p<0.05$), so analysis results obtained with the Greenhouse-Geisser correction are presented.

Table 4. Variance Analysis Results for Mean LSAS Pretest, Posttest and Follow-up Test Points for Experiment and Control Groups

Source of variance	Total squares	Degree of freedom	Mean squares	F	p	η_p^2
Between Groups		29				
Group	14873.87	1	14873.87	61.29	.000	.686
Error	6794.57	28	242.66			
Within groups						
Measurement (Pre-Post- follow-up)	11612.02	1.507	7707.94	102.45	.000	.785
Group * Measurement	10727.08	1.507	7120.53	94.64	.000	.772
Error	3173.55	42.182	75.23			

η_p^2 : partial eta square, $p < .05$

As seen in Table 4, the results of variance analysis of the points obtained on the LSAS determined the group effect caused significant difference ($F(1,28)= 61.29, p<.05$). Stated differently, without differentiating between pretest, posttest and follow-up measures in experimental and control groups, there was a significant level of difference identified in the mean points on the LSAS. Without differentiating the groups, the basic effect of measurement on the difference between mean points obtained from the pretest, posttest and follow-up test completed at different times appears to be significant ($F(1.507,42.182)= 102.45, p<.05$). Additionally, when investigating whether the variation between mean pretest, posttest and follow-up test LSAS points were different or not in the experiment and control groups, the group*measure common effect was identified to be significant ($F(1.507,42.182)= 94.64, p<.05$). The significant common effect shows the implemented program was effective in reducing the social anxiety of participants in the experiment group. Additionally, the effect of intervention and time was investigated with Wilks' lambda value and the analysis results are given in Table 5.

Table 5. Liebowitz Social Anxiety Scale (LSAS) repeated measurements Anova results according to Wilks Lamda Statistics

Effect	Wilks' λ	F	Degree of freedom	P	η_p^2
Measurement	.176	63.26	2	.000	.82
Measurement * Groups	.178	62.51	2	.000	.82

Not: Wilks' λ : Wilks Lamda Statistics, F: F-test, η_p^2 : partial eta square

As seen in Table 5, when the analysis results are investigated, the LSAS for time had Wilks' $\lambda=.176, F(2,27) = 63.26; p<.001$ and appeared to show significant variation. Similarly, the measure*group interaction effect appeared to be similar (Wilks' $\lambda=.178, F(2,27) = 62.51; p<.01$). In line with the results, mean values of measures differed between themselves as well as differing between groups. With the aim of determining the source of the difference, the Bonferroni corrected multiple comparison test was performed. The obtained results are presented in Table 6.

Table 6. Two-way Comparison (Bonferroni Correction) Test Results for LSAS Pretest, Posttest and Follow-up Test Points for Experiment and Control Groups

Groups	(I) Measurement	(J) Measurement	Difference	Standard Error	p
Experimental Group	Pretest	Posttest	46.800*	4.120	.000
	Pretest	Follow-up test	47.533*	4.128	.000
	Follow-up test	Posttest	-.733	.636	.805
Control Group	Pretest	Posttest	-.733	.714	.965
	Pretest	Follow-up test	2.533	2.023	.693
	Follow-up test	Posttest	-3.267	2.514	.644

As seen in Table 6, when the analysis results are investigated, the difference in pretest and posttest LSAS points and the difference between the pretest and follow-up test points for the solution-focused group counseling program were higher compared to the control group. Differentiation was not observed between the posttest and follow-up measures in both groups. Results are presented in Figure 1.




Figure 1.

Figure 1. Variation related to Mean LSAS Pretest, Posttest and Follow-up Test Points for Experiment and Control Groups

As seen on Figure 1, the pretests generally had points in relative proximity, while there was a sharp fall on the posttest for the solution-focused group counseling group. Finally, the solution-focused group counseling program can be said to be effective to reduce social anxiety of university students compared to the control group.

DISCUSSION

In this study to reduce the social anxiety of university students, the solution-focused group counseling program was more effective on the Liebowitz Social Anxiety Scale (LSAS) compared to a control group. This efficacy was concluded to continue during follow-up measures completed three months after the sessions ended. This result may be interpreted as showing the solution-focused group counseling program developed in this study was effective to reduce the social anxiety of university students.

This study was conducted to examine the effect of solution-focused group counseling on the social anxiety of university students. At the end of the sessions, the students evaluated this counseling process very positively. They also stated that they reduced their anxiety and fear in social situations compared to the past. The result obtained from this study is consistent with the results on social anxiety disorders based on other psychological counseling approaches other than solution-focused

group counseling (Beidel, Turner & Morris, 2000; Stangier, Heidenreich, Peitz, Lauterbach & Clark, 2003; Nedim-Bal & Öner, 2014; Palancı, 2004; Mörtberg, Karlsson, Fyring & Sundin, 2006; Zaboski, Joyce-Beaulieu, Kranzler, McNamara, Gayle & MacInnes, 2019). In this context, it can be stated that the participation of the clients in the counseling process has a positive effect on social anxiety.

As a result of literature review related to social anxiety issues in Turkey, of studies examining the effectiveness of group counseling based on solution-oriented approaches to solutions it was found to be inadequate. This study result supports the result of the study conducted by Ateş and Gençdoğan (2017). It is emphasized by De Shazer (1985) and De Shazer and Berg (1997) that solution-oriented counseling is a powerful therapeutic approach that emphasizes the resources people have and how these resources can be applied to a positive change process. In this context, with solution-oriented counseling, individuals will be able to create ways to achieve positive adaptation by using their resources when they encounter negative conditions. In addition, the result obtained from this study supports the results of psychological counseling with a solution-oriented group conducted abroad to reduce social anxiety (Bajesh, 2015; Elemo, 2019; Esmail, Alireza, Khadije, Esmail & Shima, 2019; George, 2008; Mahdiyar, Dadfarnia, Hadianfard & Rahimi, 2019). In this context, the sessions being based on the solution-focused counseling approach, focusing more on solutions to the participants' problems, and emphasizing their strong and positive aspects more than their weak points are thought to be effective on the results obtained in this research. In addition to these, it is seen that there are studies in the literature that have determined that psychological counseling with a solution-focused group is effective in dealing with various problems in life (Ateş, 2015; Ateş, 2016; Baratian, Salimi, Moghim, Shakarami, & Davarniya, 2016; Javid, Ahmadi, Mirzai & Atghaei, 2019; Mohseni Takalu, Hosseini & Khankeh, 2017; Roeden, Maaskant & Curfs, 2013; Rose & Ishak, 2019; Sağar, 2020; Sarı & Günaydın, 2016). Accordingly, the solution-focused group counseling program may have assisted university students in undergoing significant experiences and variations in terms of reducing social anxiety and discovering coping strategies. By developing resilience to social anxiety, it may have enabled them to organize their daily lives and to explore their power sources in more depth. Their own reorganization and taking action may have contributed to assessments of themselves with a positive perspective and reduced social anxiety.

In conclusion, in this study to reduce the social anxiety of university students, the solution-focused counseling program was revealed to be more effective compared to a control group. However, limiting aspects of the research are that it was completed only with university students, is not longitudinal and only compared with a control group. Additionally, just as group dynamics may have been effective on the program, they may be considered a limiting aspect. Programs to reduce social anxiety based on other psychological counseling approaches may be developed and compared with this solution-focused group counseling program to reduce social anxiety.

REFERENCES

- Abeditehrani, H., Dijk, C., Toghchi, M. S., & Arntz, A. (2020). Integrating cognitive behavioral group therapy and psychodrama for social anxiety disorder: An intervention description and an uncontrolled pilot trial. *Clinical Psychology in Europe*, 2(1), 1-21. doi: <https://doi.org/10.32872/cpe.v2i1.2693>.
- American Psychiatric Association, DSM-5.(2013). *Diagnostic and statistical manual of mental disorders: DSM-5™* (5th ed.). Washington, DC.
- Ateş, B. (2015). Üniversite öğrencilerinin güvengenlik becerileri üzerinde çözüm odaklı grupla psikolojik danışmanın etkisi. *Mersin Üniversitesi Eğitim Fakültesi Dergisi*, 11(3), 832-841., Doi: 10.17860/efd.84626

- Ateş, B. (2016). The effect of solution focused brief group counseling upon the perceived social competences of teenagers. *Journal of Education and Training Studies*, 4(7), 28-36., Doi: 10.11114/jets.v4i7.1443.
- Ateş, B. & Gençdoğan, B. (2017). Üniversite öğrencilerinin sosyal fobi ile başa çıkmalarında çözüm odaklı grupla psikolojik danışmanın etkisinin incelenmesi. *İnönü Üniversitesi Eğitim Fakültesi Dergisi*, 18(1), 188-203. doi: 10.17679/inuefd.306513.
- Aydın, A., Sütçü -Tekinsav, S. & Sorias, O. (2010). Ergenlerde sosyal anksiyete belirtilerini azaltmaya yönelik bilisel-davranışçı bir grup terapisi programının etkililiğinin değerlendirilmesi. *Türk Psikiyatri Dergisi*, 21(1),25-36.
- Baijesh, A. R. (2015). Solution focused brief therapy for social anxiety disorder-a pilot study. *International Journal of Psychology and Education*, 2 (7), 8-12.
- Baratian, A., Salimi, A., Moghim, F., Shakarami, M., & Davarniya, R. (2016). The effect of solution-focused brief therapy (SFBT) on increasing self-esteem in female students. *Iranian Journal of Nursing Research*, 11(4), 57-65.
- Baygül, Ş. (2015). *Çözüm odaklı kısa süreli aile danışmanlığı kuramına dayalı olarak geliştirilen evlilik danışmanlığı programının çiftlerin evlilik uyumlarını arttırmadaki etkisi*. Yayımlanmamış Yüksek Lisans Tezi. Muğla Sıtkı Kocaman Üniversitesi. Eğitim Bilimleri Enstitüsü, Muğla.
- Beidel, D. C., Turner, S. M. & Morris, T. L. (2000). Behavioral treatment of childhood social phobia. *Journal of Consulting and Clinical Psychology*, 68(6), 1072-1080.
- Bolsoni-Silva, A. T., & Loureiro, S. R. (2014). The role of social skills in social anxiety of university students. *Paidéia (Ribeirão Preto)*, 24(58), 223-232. doi:10.1590/1982-43272458201410.
- Büyüköztürk, Ş. (2011). *Sosyal bilimler için veri analizi el kitabı*. (15. Baskı). Ankara: Pegem A Yayıncılık.
- De Jong, P., & Berg, I. K. (1998). *Interviewing for solutions*. Pacific Grove CA: Brooks/ Cole.
- De Shazer, S. (1985). *Keys to solution in brief therapy*. New York: W.W. Norton & Company Inc.
- De Shazer, S., & Berg I. K. (1997). 'What works?' Remarks on research aspects of solution-focused brief therapy. *Journal of Family Therapy*, 19, 121-124.
- Deci, E. L., & Ryan, R. M. (Eds.). (2004). *Handbook of self-determination research*. University Rochester Press.
- Doğan, S. (1999). Çözüm odaklı kısa süreli terapi: Kuramsal bir inceleme. *Türk Psikolojik Danışma ve Rehberlik Dergisi*, 2(12), 28-38.
- Elemo, A. S. (2019). *The effect of solution-focused and social skills training groups on social anxiety and coping strategies of ethiopian university students*. Unpublished doctoral dissertation. Anadolu University. Graduate School of Educational Sciences. Eskişehir.
- Esmail, B. Alireza, B., Khadije, F., Esmail, Z., & Shima, E. (2019). The effectiveness of solution-focused brief therapy on reduction social anxiety in children and adolescents with stuttering. *Research in Clinical Psychology and Counseling (Studies in Education & Psychology)*, 9(1), 76-92.

- Fogarty, C., Hevey, D., & McCarthy, O. (2019). Effectiveness of cognitive behavioural group therapy for social anxiety disorder: Long-term benefits and aftercare. *Behavioural and Cognitive Psychotherapy*, 1–13. doi:10.1017/s1352465819000079.
- George, M. C. (2008). Solution-focused therapy: Strength-based counseling for children with social phobia. *Journal of Humanistic Counseling, Education and Development*, 47, 144-156.
- Gil, P. J. M., Carrillo, F. X. M., & Meca, J. S. (2001). Effectiveness of cognitive-behavioural treatment in social phobia: A meta-analytic review. *Psychology in Spain*, 5(1), 17–25.
- Heppner, P. P., Wampold, B. E. & Kivlighan, D. M. (2008). Gruplar arası ve denekler içi desenler (Totan, T., Çev.), *Psikolojik Danışmada Araştırma Yöntemleri*, (D. M. Siyez, Çev. Ed.). Ankara: Mentis.
- Javid, N., Ahmadi, A., Mirzai, M., & Atghaei, M. (2019). Effectiveness of solution-focused group counseling on the mental health of midwifery students. *Revista Brasileira de Ginecologia e Obstetricia*, 41(8), 500-507. doi: https://doi.org/10.1055/s-0039-1693741.
- Kim J. S., & Franklin C. (2009). *Solution-focused brief therapy in schools: A review of the outcome literature. Children and Youth Services Review*, 31(4), 464-470.
- Mahdiyari M., Dadfarnia S. H., Hadianfard H., & Rahimi C. H. (2019). Investigating the effectiveness of rational-emotional therapy and solution focused group therapy on social anxiety disorder in the students of Shiraz University. *Shenakht Journal of Psychology and Psychiatry*, 6(3), 118-128.
- Mayo-Wilson, E., Dias, S., Mavranzouli, I., Kew, K., Clark, D. M., Ades, A. E., & Pilling, S. (2014). Psychological and pharmacological interventions for social anxiety disorder in adults: A systematic review and network meta-analysis. *The Lancet Psychiatry*, 1(5), 368–376. doi:10.1016/s2215-0366(14)70329-3.
- Mercan, Ç. S. (2007). *Bilişsel-davranışçı yaklaşımla bütünleştirilmiş sosyal beceri eğitiminin ergenlerin sosyal kaygı düzeyine etkisi*. Yayınlanmamış doktora tezi. İstanbul Üniversitesi Sosyal Bilimler Enstitüsü, İstanbul.
- Mohseni Takalu, M. T., Hosseini, S. A., & Khankeh, H. (2017). Effectiveness solution-focused therapy group, to reduce stress, anxiety and depression in caregivers of patients with multiple sclerosis. *Modern Care Journal*, 14(4), 1-6. doi: 10.5812/modernc.10696.
- Mörtberg, E., Karlsson, A., Fyring, C., & Sundin, Ö. (2006). Intensive cognitive-behavioral group treatment (CBGT) of social phobia: A randomized controlled study. *Journal of Anxiety Disorders*, 20(5), 646–660. doi:10.1016/j.janxdis.2005.07.005.
- Mustafa, R. B., Hamdan-Mansour, A. M., Hijazeen, J. K., Abed, H. S., Abdallah, F. W., El-Haija, H. M. A., & Omari, H. (2014). Social phobia among university students in Jordan. *Life Science Journal*, 11(2), 93-98.
- Nedim-Bal, P. & Öner, M. (2014). Sosyal kaygı ile başa çıkma psiko eğitim programının ortaokul öğrencileri üzerindeki etkisi. *Eğitim Bilimleri Araştırmaları Dergisi*, 4(1), 335-348. http://ebad-jesr.com/.
- Palancı. M. (2004, Temmuz). Üniversite öğrencilerinin sosyal kaygı sorunlarını açıklamaya yönelik bir model çalışması. XIII. Ulusal Eğitim Bilimleri Kurultayında sunulan bildiri, İnönü Üniversitesi Eğitim Fakültesi, Malatya.

- Rapee R. M. (1995). *Social Phobia*. New York 1995.
- Roeden, J. M., Maaskant, M. A., & Curfs, L. M. G. (2013). Processes and effects of solution-focused brief therapy in people with intellectual disabilities: A controlled study. *Journal of Intellectual Disability Research*, 58(4), 307–320. doi:10.1111/jir.12038.
- Rose, N. N., & Ishak, A. S. (2019). Emotional coping through solution focused counselling therapy: Case study of psychological distressed women. *Asian Social Work Journal (ASWJ)*, 4(3), 23-34. doi:10.47405/aswj.v4i3.103.
- Sağar, M.E. (2020). *Üniversite öğrencilerinin sağlıklı internet kullanımını arttırmaya yönelik çözüm odaklı grupla psikolojik danışma programı ile grup rehberliği programının etkililiğinin incelenmesi*. Yayınlanmamış doktora tezi. Eskişehir Osmangazi Üniversitesi. Eğitim Bilimleri Enstitüsü, Eskişehir.
- Sarı, E., & Günaydın, N. (2016). Çözüm odaklı depresyon ile başa çıkma eğitiminin depresyon belirtileri üzerindeki etkileri. *Anadolu Psikiyatri Dergisi*, 17(5), 369-375.
- Simon, J., & Berg, I. K. (1997). *Solution-focused brief therapy with long-term problems*. *Directions in Rehabilitation Counseling*, 10, 117-127.
- Sklare, G. B. (1997). *Brief counseling that works: A solution-focused approach for school counselors*. California: Corwin Pres, Inc.
- Soykan, Ç. Devrimci Özgüven, H., & Gençöz, T. (2003). Liebowitz social anxiety scale: The Turkish version. *Psychological Reports*, 93(3), 1059-1069.
- Stangier, U., Heidenreich, T., Peitz, M., Lauterbach, W., & Clark, D. M. (2003). Cognitive therapy for social phobia: Individual versus group treatment. *Behaviour Research and Therapy*, 41(9), 991–1007. doi:10.1016/s0005-7967(02)00176-6.
- Tillfors, M., Andersson, G., Ekselius, L., Furmark, T., Lewenhaupt, S., Karlsson, A., & Carlbring, P. (2011). A randomized trial of internet-delivered treatment for social anxiety disorder in high school students. *Cognitive Behaviour Therapy*, 40(2), 147–157. doi:10.1080/16506073.2011.555486.
- Wintre, M. G., & Yaffe, M. (2000). First-year students' adjustment to university life as a function of relationships with parents. *Journal of Adolescent Research*, 15(1), 9–37. doi:10.1177/0743558400151002.
- Zaboski, B. A., Joyce-Beaulieu, D., Kranzler, J. H., McNamara, J. P., Gayle, C., & MacInnes, J. (2019). Group exposure and response prevention for college students with social anxiety: A randomized clinical trial. *Journal of Clinical Psychology*. 75(5), 1-19. doi:10.1002/jclp.22792.